

Chrysosplenium oppositifolium

Habitus and growth type

Plant height [m]: **0.09**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **23.35**

Fruit, seed and dispersal

Seed mass [mg]: **0.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.2**

Temperature indicator value: **5**

Moisture indicator value: **9.3**

Reaction indicator value: **5.4**

Nutrient indicator value: **5.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.36**

Disturbance frequency (herb layer): **1.43**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.04**

Grazing pressure: **0.14**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OA \(MON\) Montio-Cardaminetea](#)

Broad habitat

Occurrence in broad habitats: **Spring, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **2**

Continentality amplitude: **2**