

Impatiens glandulifera

Habitus and growth type

Plant height [m]: **1.56**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **38.3**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **8.57**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.6**

Temperature indicator value: **6.2**

Moisture indicator value: **7.6**

Reaction indicator value: **6.5**

Nutrient indicator value: **7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.52**

Disturbance frequency (herb layer): **1.3**
Disturbance severity: **0.6**
Disturbance severity (herb layer): **0.42**
Mowing frequency: **0.24**
Grazing pressure: **0.17**
Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [HA \(POP\) *Alno glutinosae-Populetea albae*](#), [QG \(EPI\) *Epilobietea angustifolii*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T11 Temperate Salix and Populus riparian forest](#)

Constant species of EUNIS habitats: [T11 Temperate Salix and Populus riparian forest](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **3**
Continentality amplitude: **3**