

Inula conyzae

Habitus and growth type

Plant height [m]: **0.66**

Life span: **Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **16.38**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.25**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.3**

Temperature indicator value: **5.9**

Moisture indicator value: **4**

Reaction indicator value: **7.2**

Nutrient indicator value: **3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.23**

Disturbance frequency (herb layer): **1.41**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.26**

Mowing frequency: **0.16**

Grazing pressure: **0.23**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) Trifolio-Geranietea sanguinei](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **4**

Continentality amplitude: **4**