

Inula montana

Habitus and growth type

Plant height [m]: **0.26**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **11.62**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.24**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **7**

Moisture indicator value: **2.3**

Reaction indicator value: **7.7**

Nutrient indicator value: **1.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.83**
Disturbance frequency (herb layer): **1.73**
Disturbance severity: **0.52**
Disturbance severity (herb layer): **0.39**
Mowing frequency: **0.37**
Grazing pressure: **0.4**
Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FN \(RUM\) Rumici-Astragaletea siculi](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#), [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#)

Constant species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#), [R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**