

Inula oculus-christi

Habitus and growth type

Plant height [m]: **0.3**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.16**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **7.4**

Moisture indicator value: **2.8**

Reaction indicator value: **8**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.09**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.4**

Grazing pressure: **0.34**

Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [KH \(DRY\) Dryidetea spinosae](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**