

# *Iris pumila*

## Habitus and growth type

Plant height [m]: **0.11**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **25.28**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **6.9**

Moisture indicator value: **2.5**

Reaction indicator value: **9**

Nutrient indicator value: **1.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.19**

Disturbance frequency (herb layer): **1.9**

Disturbance severity: **0.47**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.43**

Grazing pressure: **0.39**

Soil disturbance: **0.18**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1B Continental dry grassland \(true steppe\)](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**