

# *Jacobaea adonidifolia*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **4**

Moisture indicator value: **3**

Reaction indicator value: **4.5**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.71**

Disturbance frequency (herb layer): **1.55**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.34**

Mowing frequency: **0.23**

Grazing pressure: **0.27**

Soil disturbance: **0.15**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CH \(ULI\) \*Calluno-Ulicetea\*](#), [CL \(GER\) \*Trifolio-Geranietea sanguinei\*](#), [CT \(TRI\) \*Juncetea trifidi\*](#), [FE \(CYT\) \*Cytisetea scopario-striati\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Heathland, Scrub**