Jacobaea carniolica

Habitus and growth type

Plant height [m]: **0.12**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.76**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **1.5**
Moisture indicator value: **5**
Reaction indicator value: **1**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.33**
Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.29**
Disturbance severity (herb layer): **0.27**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0**
Grazing pressure: **0.22**
Soil disturbance: **0.09**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CT (TRI) *Juncetea trifidi***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R43 Temperate acidophilous alpine grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),**
Oromediterranean scrub

Distribution

Continentality: **5**
Continentality amplitude: **1**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz