Jasminum fruticans

Habitus and growth type

Plant height [m]: **1.5**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Specific leaf area [mm^2/mg]: **14.12**

Flower

Flowering period: **January-December**

Fruit, seed and dispersal

Seed mass [mg]: **67.65**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.9**
Temperature indicator value: **8.2**
Moisture indicator value: **3.3**
Reaction indicator value: **6.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.2**
Disturbance frequency (herb layer): **1.31**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.05**
Grazing pressure: **0.27**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CB (PUB) *Quercetea pubescentis*, FA (QUI) *Quercetea ilicis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3D Mediterranean Cupressaceae forest**
Constant species of EUNIS habitats: **T3D Mediterranean Cupressaceae forest**

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz