

# *Juncus capitatus*

## Habitus and growth type

Plant height [m]: **0.08**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.01**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.1**

Nutrient indicator value: **2**

Salinity indicator value: **0.4**

### Disturbance indicator values

Disturbance frequency: **1.92**

Disturbance frequency (herb layer): **2.06**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.51**

Mowing frequency: **0.2**

Grazing pressure: **0.23**

Soil disturbance: **0.31**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation](#), [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Spring, Wetland, Grassland (non-alpine, non-saline), Saline vegetation, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **4**

Continentality amplitude: **4**