

# *Jurinea stoechadifolia*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **5.4**

Moisture indicator value: **4.4**

Disturbance indicator values

Disturbance frequency: **1.76**

Disturbance frequency (herb layer): **1.76**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.42**

Grazing pressure: **0.4**

Soil disturbance: **0.17**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#)

Constant species of EUNIS habitats: [R15 Continental dry rocky steppic grassland](#)

[and dwarf scrub on chalk outcrops](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**