

# *Koeleria crassipes*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **6**

Moisture indicator value: **2**

Reaction indicator value: **5**

Nutrient indicator value: **1**

### Disturbance indicator values

Disturbance frequency: **0.76**

Disturbance frequency (herb layer): **1.43**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.09**

Grazing pressure: **0.32**

Soil disturbance: **0.13**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FL \(IND\) \*Festucetea indigestae\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R1N Open Iberian supramediterranean dry acid and neutral grassland](#)

Constant species of EUNIS habitats: [R1N Open Iberian supramediterranean dry acid and neutral grassland](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**