

# *Koeleria eriostachya*

## Habitus and growth type

Plant height [m]: **0.39**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **June-August**

## Fruit, seed and dispersal

Seed mass [mg]: **0.47**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Moisture indicator value: **3**

Reaction indicator value: **7**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.9**

Disturbance frequency (herb layer): **1.36**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.08**

Grazing pressure: **0.24**

Soil disturbance: **0.08**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CU \(SES\) \*Elyno-Seslerietea\*](#), [DA \(FES\) \*Festuco-Brometea\*](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),  
Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**