

Koeleria lobata

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **7.1**

Moisture indicator value: **4**

Reaction indicator value: **8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.19**

Disturbance frequency (herb layer): **1.9**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.2**

Grazing pressure: **0.37**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FN \(RUM\) Rumici aetnensis-Astragaletea siculi](#), [FR \(DAP\) Daphno-Festucetea](#), [KH \(DRY\) Drypidetea spinosae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#), [R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#)

Constant species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#), [R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Sparsely vegetated (incl. rock and scree)**