

# ***Lactuca perennis***

## **Habitus and growth type**

Plant height [m]: **0.43**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **23.47**

## **Flower**

Flowering period: **March-July**

## **Fruit, seed and dispersal**

Seed mass [mg]: **1.37**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **9**

Temperature indicator value: **6.9**

Moisture indicator value: **2.2**

Reaction indicator value: **8**

Nutrient indicator value: **2.3**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.58**

Disturbance frequency (herb layer): **1.49**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.2**

Grazing pressure: **0.29**

Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FP \(ONO\) Festuco hystricis-Ononidetea striatae](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [U72 Unvegetated or sparsely vegetated gravel bar in the Mediterranean region](#)

### Broad habitat

Occurrence in broad habitats: **Coastal cliff, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **5**

Continentality amplitude: **3**