

Lathyrus pisiformis

Habitus and growth type

Plant height [m]: **0.75**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

Fruit, seed and dispersal

Seed mass [mg]: **13.98**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.7**

Temperature indicator value: **5.2**

Moisture indicator value: **4.5**

Reaction indicator value: **8**

Nutrient indicator value: **4.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.44**

Disturbance frequency (herb layer): **1.58**

Disturbance severity: **0.59**

Disturbance severity (herb layer): **0.36**

Mowing frequency: **0.37**

Grazing pressure: **0.24**

Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BC \(BRA\) *Brachypodio pinnati-Betuletea pendulae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H *Larix light taiga*](#)

Constant species of EUNIS habitats: [T3H *Larix light taiga*](#)

Broad habitat

Occurrence in broad habitats: **Forest**