

Laurus nobilis

Habitus and growth type

Plant height [m]: **9.62**

Life span: **Perennial**

Life form: **Phanerophyte, Tree**

Leaf

Specific leaf area [mm²/mg]: **5.1**

Flower

Flowering period: **February-April**

Fruit, seed and dispersal

Seed mass [mg]: **615.3**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **8.6**

Moisture indicator value: **5.8**

Reaction indicator value: **5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.08**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.15**

Mowing frequency: **0**

Grazing pressure: **0.23**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) *Quercetea ilicis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T22 Mainland laurophyllous forest](#)

Constant species of EUNIS habitats: [T22 Mainland laurophyllous forest](#)

Dominant species of EUNIS habitats: [T22 Mainland laurophyllous forest](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**