

# *Leopoldia comosa*

## Habitus and growth type

Plant height [m]: **0.47**

Life span: **Perennial**

Life form: **Geophyte**

Spinescence: **not spinescent**

## Flower

Flowering period: **March-June**

## Fruit, seed and dispersal

Seed mass [mg]: **5.19**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **7.4**

Moisture indicator value: **3.4**

Reaction indicator value: **6.7**

Nutrient indicator value: **4.8**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.6**

Disturbance frequency (herb layer): **1.66**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.48**

Mowing frequency: **0.36**

Grazing pressure: **0.26**

Soil disturbance: **0.29**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality amplitude: **8**