

Lepidium densiflorum

Habitus and growth type

Plant height [m]: **0.28**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **60.09**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.53**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **6.8**

Moisture indicator value: **3.7**

Reaction indicator value: **7.4**

Nutrient indicator value: **5.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **1.96**

Disturbance severity: **0.76**

Disturbance severity (herb layer): **0.74**

Mowing frequency: **0.95**

Grazing pressure: **0.16**

Soil disturbance: **0.54**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QD \(DIG\) *Digitario sanguinalis-Eragrostietea minoris*](#)

Broad habitat

Occurrence in broad habitats: **Synanthropic**

Distribution

Continentality: 7

Continentality amplitude: 6