

Limodorum abortivum

Habitus and growth type

Plant height [m]: **0.49**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **0**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **full mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.2**

Moisture indicator value: **4.3**

Reaction indicator value: **7.7**

Nutrient indicator value: **3.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.06**

Disturbance frequency (herb layer): **1.35**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.14**

Mowing frequency: **0.03**

Grazing pressure: **0.25**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

Distribution

Continentality amplitude: **5**