# *Limoniastrum monopetalum*

## Habitus and growth type

Plant height [m]: **0.75**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

## Flower

Flowering period: **May-August**

## Fruit, seed and dispersal

Seed mass [mg]: **9.36**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Saline**

### Ellenberg-type indicator values

Light indicator value: **8.7**
Temperature indicator value: **10.3**
Moisture indicator value: **8**
Reaction indicator value: **8.7**
Nutrient indicator value: **6.7**
Salinity indicator value: **8.3**

### Disturbance indicator values

Disturbance frequency: **1.54**
Disturbance frequency (herb layer): **2.17**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.45**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0**
Grazing pressure: **0.19**
Soil disturbance: **0.26**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **MG (SAL) *Salicornietea fruticosae***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **MA253 Mediterranean mid-low saltmarsh**

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Saline vegetation**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*