Limonium cordatum

Habitus and growth type

Plant height [m]: **0.2**
Life span: **Perennial**
Life form: **Chamaephyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **9.5**
Moisture indicator value: **4.5**
Reaction indicator value: **7**
Nutrient indicator value: **4.5**
Salinity indicator value: **7**

Disturbance indicator values

Disturbance frequency: **0.54**
Disturbance frequency (herb layer): **0.54**
Disturbance severity: **0.12**
Disturbance severity (herb layer): **0.1**
Mowing frequency: **0**
Grazing pressure: **0.01**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.01**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JB (CRI) *Crithmo-Staticetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **N32 Mediterranean and Black Sea rocky sea cliff and shore**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz