

Limonium graecum

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **11.5**

Reaction indicator value: **9**

Nutrient indicator value: **6.5**

Salinity indicator value: **5.5**

Disturbance indicator values

Disturbance frequency: **0.76**

Disturbance frequency (herb layer): **0.79**

Disturbance severity: **0.34**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.07**

Grazing pressure: **0.1**

Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JB \(CRI\) Crithmo-Staticetea](#), [MG \(SAL\) Salicornietea fruticosae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N32 Mediterranean and Black Sea rocky sea cliff and shore](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Saline vegetation**