

Linaria spartea

Habitus and growth type

Plant height [m]: **0.38**

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.04**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7**

Moisture indicator value: **1**

Reaction indicator value: **4**

Nutrient indicator value: **4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.99**

Disturbance frequency (herb layer): **1.85**

Disturbance severity: **0.44**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.55**

Grazing pressure: **0.29**

Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) *Helianthemetea guttati*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1R Mediterranean to Atlantic open, dry, acid and neutral grassland](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

Distribution

Continentality: **2**

Continentality amplitude: **2**