

Lindernia dubia

Habitus and growth type

Plant height [m]: **0.14**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**

Dispersal mode: **Anthropochory, Hydrochory**

Dispersal distance class: **7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **7**

Moisture indicator value: **8.6**

Reaction indicator value: **6.6**

Nutrient indicator value: **6.7**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.88**

Disturbance frequency (herb layer): **1.88**

Disturbance severity: **0.82**

Disturbance severity (herb layer): **0.82**

Mowing frequency: **0.04**

Grazing pressure: **0.16**

Soil disturbance: **0.46**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#), [QJ \(ORY\) Oryzetea sativae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V14 Inundated or inundatable cropland, including rice fields](#)

Constant species of EUNIS habitats: [V14 Inundated or inundatable cropland, including rice fields](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Synanthropic**

Distribution

Continentality amplitude: **5**