

# *Lindernia procumbens*

## Habitus and growth type

Plant height [m]: **0.08**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **June-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.01**

Dispersal mode: **Hydrochory**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.9**

Temperature indicator value: **7.1**

Moisture indicator value: **8.2**

Reaction indicator value: **7.3**

Nutrient indicator value: **6.5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.83**

Disturbance frequency (herb layer): **1.83**

Disturbance severity: **0.71**

Disturbance severity (herb layer): **0.71**

Mowing frequency: **0.04**

Grazing pressure: **0.16**

Soil disturbance: **0.38**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V14 Inundated or inundatable cropland, including rice fields](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Synanthropic**

## **Distribution**

Continentality: **7**

Continentality amplitude: **8**