Lindernia procumbens

Habitus and growth type

Plant height [m]: **0.08**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**
Dispersal mode: **Hydrochory**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.9**
Temperature indicator value: **7.1**
Moisture indicator value: **8.2**
Reaction indicator value: **7.3**
Nutrient indicator value: **6.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.83**
Disturbance frequency (herb layer): **1.83**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.71**
Mowing frequency: **0.04**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.16**
Soil disturbance: **0.38**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OC (ISO) *Isoëto-Nanojuncetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **V14 Inundated or inundatable cropland, including rice fields**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Synanthropic**

Distribution

Continentality: **7**
Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz