

Linum bienne

Habitus and growth type

Plant height [m]: **0.33**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **1.27**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**

Moisture indicator value: **3.6**

Reaction indicator value: **6.5**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.39**

Disturbance frequency (herb layer): **1.96**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.54**

Mowing frequency: **0.55**

Grazing pressure: **0.28**

Soil disturbance: **0.3**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) *Molinio-Arrhenatheretea*](#),
[FG \(SAC\) *Stipo giganteae-Agrostietea castellanae*](#)

EUNIS habitat

Constant species of EUNIS habitats: [S34 *Balkan-Anatolian submontane genistoid scrub*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**