

# *Linum tauricum*

## Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **1.29**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5.8**

Moisture indicator value: **4.8**

Disturbance indicator values

Disturbance frequency: **1.7**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.6**

Grazing pressure: **0.38**

Soil disturbance: **0.18**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

Constant species of EUNIS habitats: [R17 Heavy-metal dry grassland of the Balkans](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely**

**vegetated (incl. rock and scree)**