

Lomelosia cretica

Habitus and growth type

Plant height [m]: **0.7**

Life span: **Perennial**

Life form: **Chamaephyte**

Flower

Flowering period: **March-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **10**

Moisture indicator value: **1**

Reaction indicator value: **8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.93**

Disturbance frequency (herb layer): **1.36**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.39**

Mowing frequency: **0.21**

Grazing pressure: **0.41**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) *Asplenietea trichomanis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S54 Thermomediterranean arid scrub](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Scrub, Sparsely vegetated (incl. rock and scree)**