

# *Lonicera tatarica*

## Habitus and growth type

Plant height [m]: **2.13**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

## Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**

Leaf phenology (woody plants only): **Winter deciduous**

## Fruit, seed and dispersal

Seed mass [mg]: **3.18**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **6.5**

Temperature indicator value: **6.8**

Moisture indicator value: **6**

Disturbance indicator values

Disturbance frequency: **0.29**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.67**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.1**

Grazing pressure: **0.19**

Soil disturbance: **0.16**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [HA \(POP\) Alno glutinosae-Populetea albae](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest**