Lotus borbasii

Habitus and growth type

Plant height [m]: **0.25**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **1.24**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **7.5**
Moisture indicator value: **2.5**
Reaction indicator value: **8.5**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.47**
Disturbance frequency (herb layer): **1.96**
Disturbance severity: **0.48**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.68**
Grazing pressure: **0.32**
Soil disturbance: **0.14**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz