

# *Lotus hispidus*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.44**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **6**

Moisture indicator value: **1**

Reaction indicator value: **2**

Nutrient indicator value: **1**

Salinity indicator value: **0.1**

### Disturbance indicator values

Disturbance frequency: **1.73**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.28**

Grazing pressure: **0.25**

Soil disturbance: **0.28**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [Q62 Periodically exposed shore with stable,](#)

mesotrophic sediments with pioneer or ephemeral vegetation

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline)**