Lycium schweinfurthii

Habitus and growth type

Life span: **Perennial**
Life form: **Phanerophyte, Shrub**
Spinescence: **spinescent**

Fruit, seed and dispersal

Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **12**
Moisture indicator value: **1**
Reaction indicator value: **8**
Nutrient indicator value: **8**
Salinity indicator value: **3**

Disturbance indicator values

Disturbance frequency: **0.91**
Disturbance frequency (herb layer): **1.6**
Disturbance severity: **0.59**
Disturbance severity (herb layer): **0.44**
Mowing frequency: **0.09**
Grazing pressure: **0.27**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of phytosociological classes: **FK (PEG) *Pegano harmalae-Salsoletea vermiculatae***

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz