Lythrum tribracteatum

Habitus and growth type

Plant height [m]: **0.18**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.02**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **7.6**
Moisture indicator value: **7.5**
Nutrient indicator value: **3.8**
Salinity indicator value: **0.6**

Disturbance indicator values

Disturbance frequency: **0.98**
Disturbance frequency (herb layer): **1.79**
Disturbance severity: **0.62**
Disturbance severity (herb layer): **0.56**
Mowing frequency: **0.07**
Grazing pressure: **0.21**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OC (ISO) *Isoëto-Nanojuncetea***

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz