

Melampyrum arvense

Habitus and growth type

Plant height [m]: **0.32**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **28.26**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **13.82**

Dispersal mode: **Myrmecochory, Anthropochory**

Dispersal distance class: **3, 7**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **6.4**

Moisture indicator value: **3.8**

Reaction indicator value: **8**

Nutrient indicator value: **3.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.17**

Disturbance frequency (herb layer): **1.81**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.49**
Mowing frequency: **0.67**
Grazing pressure: **0.29**
Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [QA \(PAR\) Papaveretea rhoeadis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

Distribution

Continentality: **6**
Continentality amplitude: **5**