

Melica nutans

Habitus and growth type

Plant height [m]: **0.34**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **28.73**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **3.01**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **4.3**

Temperature indicator value: **5.3**

Moisture indicator value: **4.6**

Reaction indicator value: **6.6**

Nutrient indicator value: **3.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.05**

Disturbance frequency (herb layer): **1.29**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0.02**
Grazing pressure: **0.19**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BC \(BRA\) *Brachypodio pinnati-Betuletea pendulae*](#), [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3F Dark taiga](#), [T3H Larix light taiga](#)

Constant species of EUNIS habitats: [T17 Fagus forest on non-acid soils](#), [T1E Carpinus and Quercus mesic deciduous forest](#), [T1F Ravine forest](#), [T31 Temperate mountain Picea forest](#), [T3F Dark taiga](#), [T3H Larix light taiga](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Synanthropic**

Distribution

Continentality: **6**
Continentality amplitude: **5**