

Meniocus linifolius

Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.73**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **5.8**

Moisture indicator value: **3.5**

Disturbance indicator values

Disturbance frequency: **2.11**

Disturbance frequency (herb layer): **2.07**

Disturbance severity: **0.44**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.28**

Grazing pressure: **0.41**

Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FJ \(TRA\) *Stipo-Trachynietea distachyae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R15 Continental dry rocky steppic grassland](#)

[and dwarf scrub on chalk outcrops, R1C Desert steppe, S67 Aralo-Caspian semi-desert](#)

Constant species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops, R1C Desert steppe, S67 Aralo-Caspian semi-desert](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Semi-desert**