

# *Mibora minima*

## Habitus and growth type

Plant height [m]: **0.08**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **February-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.13**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **8.3**

Moisture indicator value: **2**

Nutrient indicator value: **2.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **2**

Disturbance frequency (herb layer): **2.18**

Disturbance severity: **0.43**

Disturbance severity (herb layer): **0.43**

Mowing frequency: **0.35**

Grazing pressure: **0.29**

Soil disturbance: **0.46**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CJ \(COR\) \*Koelerio-Corynephoretea canescentis\*](#), [FI \(TUB\) \*Helianthemetea guttati\*](#), [QA \(PAR\) \*Papaveretea rhoeadis\*](#)

### Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality: **2**

Continentality amplitude: **2**