# *Minuartia attica*

## Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7**
Moisture indicator value: **4**
Reaction indicator value: **8**
Nutrient indicator value: **6**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.02**
Disturbance frequency (herb layer): **1.61**
Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.35**
Mowing frequency: **0.09**
Grazing pressure: **0.32**
Soil disturbance: **0.14**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **FR (DAP) *Daphno-Festucetea***

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## EUNIS habitat

Diagnostic species of EUNIS habitats: **R1K Balkan and Anatolian oromediterranean dry grassland**, **S75 Eastern Mediterranean mountain hedgehog-heath**

Constant species of EUNIS habitats: **R1K Balkan and Anatolian oromediterranean dry grassland**, **S75 Eastern Mediterranean mountain hedgehog-heath**

## Broad habitat

Occurrence in broad habitats: **Oromediterranean scrub**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*