Minuartia hamata

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.4**
Temperature indicator value: **6.8**
Moisture indicator value: **3.1**
Reaction indicator value: **7.1**
Nutrient indicator value: **2.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.89**
Disturbance frequency (herb layer): **1.97**
Disturbance severity: **0.6**
Disturbance severity (herb layer): **0.54**
Mowing frequency: **0.34**
Grazing pressure: **0.33**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FI (TUB) *Helianthemetea guttati***, **FR (DAP) *Daphno-Festucetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R17 Heavy-metal dry grassland of the**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Balkans, T3D Mediterranean Cupressaceae forest

Constant species of EUNIS habitats: **T3D Mediterranean Cupressaceae forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz