

Moneses uniflora

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Chamaephyte**

Leaf

Specific leaf area [mm²/mg]: **19.9**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **3.9**

Temperature indicator value: **4**

Moisture indicator value: **5.3**

Reaction indicator value: **4**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.04**

Disturbance frequency (herb layer): **1.02**

Disturbance severity: **0.66**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0.01**

Grazing pressure: **0.19**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**BA \(PIC\) Vaccinio-Piceetea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**N1F Baltic coniferous coastal dune forest**](#)

Constant species of EUNIS habitats: [**N1F Baltic coniferous coastal dune forest**](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

Distribution

Continentality: **6**

Continentality amplitude: **5**