

# *Nananthea perpusilla*

## Habitus and growth type

Plant height [m]: **0.02**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **January-June**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **10**

Moisture indicator value: **5**

Reaction indicator value: **6**

Nutrient indicator value: **1**

Salinity indicator value: **3**

### Disturbance indicator values

Disturbance frequency: **2.06**

Disturbance frequency (herb layer): **2.06**

Disturbance severity: **0.25**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.04**

Grazing pressure: **0.05**

Soil disturbance: **0.12**

### **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [OC \(ISO\) Isoëto-Nanojuncetea](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation**