

Neotinea ustulata

Habitus and growth type

Plant height [m]: **0.27**

Life span: **Perennial**

Life form: **Geophyte**

Leaf

Specific leaf area [mm²/mg]: **19.82**

Flower

Flowering period: **January-August, December**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **4.8**

Reaction indicator value: **6.7**

Nutrient indicator value: **2.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.41**

Disturbance frequency (herb layer): **1.87**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.95**

Grazing pressure: **0.27**

Soil disturbance: **0.16**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

Distribution

Continentality: **5**

Continentality amplitude: **4**