

Neottia cordata

Habitus and growth type

Plant height [m]: **0.1**

Life span: **Perennial**

Life form: **Geophyte**

Leaf

Specific leaf area [mm²/mg]: **36.51**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **2.7**

Temperature indicator value: **3.7**

Moisture indicator value: **6.7**

Reaction indicator value: **2.2**

Nutrient indicator value: **2.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.09**

Disturbance frequency (herb layer): **1.13**

Disturbance severity: **0.63**

Disturbance severity (herb layer): **0.13**

Mowing frequency: **0.01**

Grazing pressure: **0.2**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**BA \(PIC\) Vaccinio-Piceetea**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**N1F Baltic coniferous coastal dune forest**](#),
[**U35 Boreal and Arctic base-rich inland cliff**](#)

Constant species of EUNIS habitats: [**N1F Baltic coniferous coastal dune forest**](#), [**T3K Picea mire forest**](#), [**U35 Boreal and Arctic base-rich inland cliff**](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **5**