

# *Oenanthe pimpinelloides*

## Habitus and growth type

Plant height [m]: **0.57**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **2**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.7**

Temperature indicator value: **8.3**

Moisture indicator value: **6**

Reaction indicator value: **5.5**

Nutrient indicator value: **4**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **0.18**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.24**

Mowing frequency: **0.1**

Grazing pressure: **0.23**

Soil disturbance: **0.15**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) \*Molinio-Arrhenatheretea\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

Constant species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#)

### Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Scrub, Forest**