

Oenanthe pimpinelloides

Habitus and growth type

Plant height [m]: **0.57**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Seed mass [mg]: **2**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.7**

Temperature indicator value: **8.3**

Moisture indicator value: **6**

Reaction indicator value: **5.5**

Nutrient indicator value: **4**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.18**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.24**

Mowing frequency: **0.1**

Grazing pressure: **0.23**

Soil disturbance: **0.15**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S34 Balkan-Anatolian submontane genistoid scrub](#), [T1G Alnus cordata forest](#)

Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Scrub, Forest**