

# *Ochlopoa infirma*

## Habitus and growth type

Plant height [m]: **0.05**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **February-September**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7.8**

Reaction indicator value: **5.8**

Nutrient indicator value: **5.2**

Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **1.35**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.67**

Mowing frequency: **0.41**

Grazing pressure: **0.32**

Soil disturbance: **0.38**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [QE \(POL\) \*Polygono-Poetea annuae\*](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**