

# *Onobrychis viciifolia*

## Habitus and growth type

Plant height [m]: **0.38**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **18.83**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Seed mass [mg]: **19.21**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.9**

Temperature indicator value: **6.5**

Moisture indicator value: **3.8**

Reaction indicator value: **8**

Nutrient indicator value: **3.2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.58**

Disturbance frequency (herb layer): **1.91**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.49**

Mowing frequency: **1.03**

Grazing pressure: **0.26**

Soil disturbance: **0.19**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality: 7

Continentality amplitude: 7