

# *Ophrys bombyliflora*

## Habitus and growth type

Plant height [m]: **0.15**

Life span: **Perennial**

Life form: **Geophyte**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **initial or partial mycoheterotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **9.3**

Moisture indicator value: **3.7**

Reaction indicator value: **7**

Nutrient indicator value: **3**

Salinity indicator value: **0**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) \*Ononido-Rosmarinetea\*](#), [FF \(LYG\) \*Lygeo sparti-Stipetea tenacissimae\*](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**